

Law Enforcement Fatigue, Sleep, Rest and Risk Considerations

The materials provided in this correspondence are for general informational and educational purposes only and are not intended to be and should not be considered legal advice or opinions. Prior to making any policy or rule changes seek the advice of your municipal attorney.

J.A. Montgomery
CONSULTING

TIRED COPS RESEARCH FOR POLICE CHIEFS:

This is a large study that the U.S. Department of Justice prepared:

The Impact of Shift Length in Policing on Performance, Health, Quality of Life, Sleep, Fatigue, and Extra-Duty Employment

<https://www.ojp.gov/pdffiles1/nij/grants/237330.pdf>

Police Fatigue

<https://www.ojp.gov/pdffiles1/jr000248d.pdf>

Fatigue and on-duty injury among police officers: The BCOPS Study

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6311701/>

Dead Tired, Fighting fatigue (video)

<https://www.police1.com/officer-safety/articles/dead-tired-fighting-fatal-fatigue-in-law-enforcement-7NMhNEBJ7QWBYgxV/>

Police Chiefs Magazine Article

<https://www.policechiefmagazine.org/human-fatigue-in-247-operations/>

TEDx Spokane, Speech on "Tired COPS, justice and injustice, by Bryan Vila

<https://www.youtube.com/watch?v=A6eHzjEGQMo>

The Effects of Fatigue on Cognitive Performance in Police Officers and Staff During a Forward Rotating Shift Pattern

<https://reader.elsevier.com/reader/sd/pii/S2093791118301069?token=CAE30247837E14D0F0334C4814F5733F8B65BE5DFFF1B4AD25A42958240A8817CDB49F739836BBBB581C5CF838DAD6F2&originRegion=us-east-1&originCreation=20210628202114>

IACP Officer Family Wellness Guide:

<https://www.theiacp.org/resources/document/employee-and-family-wellness-guide>

-----Book: Tired Cops; The importance of Managing Police Fatigue

Police Executive Research Forum ISBN 1-878734-67-9



Published in final edited form as:

J Safety Res. 2017 February ; 60: 43–51. doi:10.1016/j.jsr.2016.11.006.

Fatigue and on-duty injury among police officers: The BCOPS study

Desta Fekedulegn^{a,*}, Cecil M. Burchfiel^a, Claudia C. Ma^a, Michael E. Andrew^a, Tara A. Hartley^a, Luenda E. Charles^a, Ja K. Gu^a, and John M. Violanti^b

^aBiostatistics and Epidemiology Branch, Health Effects Laboratory Division, National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, Morgantown, WV, USA

^bDepartment of Epidemiology and Environmental Health, School of Public Health and Health Professions, The State University of New York at Buffalo, Buffalo, NY, USA

Abstract

Introduction: Policing involves inherent physical and psychological dangers as well as occupational stressors that could lead to chronic fatigue. Although accounts of adverse events associated with police fatigue are not scarce, literature on the association between chronic fatigue and on-duty injury are limited.

Methods: Participants were officers from the Buffalo Cardio-Metabolic Occupational Police Stress (BCOPS) Study. A 10-item questionnaire was administered to assess how tired or energetic the officers generally felt irrespective of sleep hours or workload. The questionnaire consisted of five positively worded and five negatively phrased items that measured feelings of vigor/energy and tiredness, respectively. Total as well as separate scores for positive and negative items were computed by summing scores of individual items. Payroll records documenting each officer's work history were used to assess occurrence of injury. Poisson regression was used to estimate prevalence ratios (PR) of injury.

Results: Nearly 40% of officers reported feeling drained. *Overall* prevalence of *on-duty injury during the past year* was 23.9%. Injury prevalence showed a significant increasing trend across tertiles of total fatigue score: 19.6, 21.7, and 30.8% for lowest, middle and highest tertiles, respectively (trend *p*-value = 0.037). After controlling for potential confounders, a 5-unit increase in total fatigue score was associated with a 12% increase in prevalence of injury which was marginally significant (*p* = 0.075). A 5-unit increase in fatigue score of the positively worded items was associated with a 33% increase in prevalence of injury (PR = 1.33, 95% CI: 1.04–1.70, *p* = 0.022).

Conclusion: Officers who do not feel active, full of vigor, alert, or lively had a significantly higher prevalence of non-fatal work place injury compared to their counter parts.

*Corresponding author at: National Institute for Occupational Safety and Health, HELD/BEB, MS L-4050, 1095 Willowdale Rd., Morgantown, WV 26505-2888, USA. djf7@cdc.gov (D. Fekedulegn).

Declaration of interest

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Practical applications: With additional prospective evidence, workplace interventions designed to enhance level of energy may reduce feelings of tiredness and hence may prevent workplace injury.

Keywords

Non-fatal injuries; Chronic tiredness; Law enforcement; Work history

1. Introduction

Fatigue, broadly defined as “a feeling of weariness, tiredness, or lack of energy,” is a frequently cited complaint among the U.S. workforce with reported prevalence of 38% (Ricci, Chee, Lorandean, & Berger, 2007). It is an especially serious concern among police officers who are overly fatigued because of long and irregular work hours, shift work, sleep deprivation, and the inherent physical and psychosocial danger associated with the job (Vila, 2006; Vila & Kenney, 2002). Law enforcement is also one of six occupations with the highest incidence rates of nonfatal occupational injuries. The most recent data provided by U.S. Bureau of Labor Statistics (BLS) indicated that, in 2014, police and Sheriffs patrol officers had one of the highest days away from work (DAFW) nonfatal injury rates (485.8 per 10,000 full-time workers) among all occupations (107.1 per 10,000 full-time workers) and incurred the highest number of injuries among local government and second highest among state government employees (BLS, 2014). Fatigue in police officers impairs vigilance, reaction time, and performance thereby elevating the risk for fatal and non-fatal injuries to both the officers and the general public (Garbarino et al., 2007; Rajaratnam et al., 2011; Vila, 2006; Vila & Kenney, 2002; Waggoner, Grant, Van Dongen, Belenky, & Vila, 2012).

While considerable attention has been placed on the psychosocial and cardio-metabolic health of police officers (Violanti et al., 2006), scientific research on occupational injury of officers is limited, and statistics for injuries are less readily available (LaTourrette, 2011). In 2009, the National Public Safety Agenda, which is part of the National Occupational Research Agenda (NORA) for occupational safety and health research and practice in the United States, recognized that data on occupational injuries and illness among law enforcement personnel are not sufficient (NORA, 2009). Fatigue is a well-known risk factor for injury, yet the scientific literature documenting the prevalence of fatigue among police officers, particularly its association with non-fatal on-duty injury, is limited (James & Vila, 2015).

Fatigue is a latent construct that cannot be directly measured. It is argued that fatigue is best viewed as a continuum (Lewis & Wessely, 1992; Ricci et al., 2007), where at the lower end it occurs frequently and consists of acute episodes that resolve quickly following an intervention (e.g., rest, improvement of the stressor), while at the severe end it occurs less frequently but is symptomatic of a more chronic and potentially disabling conditions that cannot be quickly resolved with rest (often referred to as chronic fatigue). Chronic fatigue was defined by Barton et al. (1995) as “a general tiredness and lack of energy irrespective of whether an individual has not had enough sleep or has been working hard, which persists

even on rest days and holidays.” Although there is no standard way to assess fatigue, there are a variety of questionnaires, with high reliability and validity, which have been designed to assess fatigue in working populations (De Vries, Michielsen, & Van Heck, 2003). However, the applicability of these instruments in assessing fatigue prevalence in police officers has not been explored.

Prior studies that highlighted the significant impact of fatigue on injury and performance in police officers (James & Vila, 2015; Senjo, 2011; Vila, 2006; Violanti et al., 2012, 2013) utilized proxy indicators of fatigue (e.g., shift work, long work hours, insufficient sleep); rather than chronic fatigue assessed using one of several validated instruments. Therefore, the purpose of this research was to estimate the prevalence of chronic fatigue (assessed based on a validated instrument) and then examine its association with non-fatal workplace injury (objectively assessed using organizational work history records), among officers working in mid-sized urban police department. In our analysis, the association of interest was adjusted for demographic and lifestyle factors that were reported to affect non-fatal occupational injury in various occupational groups. These factors included age (Landen & Hendricks, 1992), gender (Berecki-Gisolf, Smith, Collie, & McClure, 2015), race/ethnicity (Hurley & Lebbon, 2012), education (Kim et al., 2014), workload (Nakata et al., 2006), physical activity (Caban-Martinez et al., 2015), and alcohol consumption (Stallones & Xiang, 2003).

2. Methods

2.1. Study population

Participants from the Buffalo Cardio-Metabolic Occupational Police Stress (BCOPS) study were used for the current analyses. The BCOPS study was a cross-sectional study aimed at investigating the associations of occupational stressors with the psychological and physiological health of police officers. The study was initiated in 2004 and a total of 710 police officers who worked with the Buffalo Police Department in New York were invited to participate in the BCOPS study; 464 (65.4%) officers agreed to participate and were examined once during the period of June 4, 2004 to October 2, 2009. No specific inclusion criteria were indicated for the study, only that participants be a sworn police officer and willing to participate. Comparisons of available demographic variables (sex, age, and police rank) showed no significant differences between participants and non-participants. A written informed consent was collected from each participant. Data collection was performed at The Center for Preventive Medicine, State University of New York at Buffalo. The study was approved by the Internal Review Board of the State University of New York at Buffalo, and the National Institute for Occupational Safety and Health (NIOSH) Institutional Review Board (IRB).

2.2. Measures and study design

Data originated from two sources. The first source was the BCOPS study where data on demographic, physical, biological, and psychosocial characteristics were collected from each participant. As part of the study, the participants filled out a questionnaire designed to assess chronic fatigue which served as the exposure variable of interest. The second source

was work history records of the BCOPS study participants obtained from the Buffalo, NY police payroll department. The work history records were used to derive occurrence of nonfatal on-duty injury which served as the outcome variable of interest.

2.3. Assessment of chronic fatigue

Chronic fatigue was assessed using a 10-item questionnaire developed by Barton et al. (1995). In the current study the questions about chronic fatigue were introduced with this statement: “The following items relate to how tired or energetic you generally feel, irrespective of whether you have had enough sleep or have been working very hard. Some people appear to suffer from permanent tiredness, even on rest days and holidays, while others seem to have limitless energy. Please indicate the degree to which the following statements apply to your own normal feelings.” The study participants were then asked to rate (score) each of the 10 items on a five-point Likert scale (5 = very much, 4 = much, 3 = somewhat, 2 = little, 1 = not at all). The questionnaire consisted of five items (I usually feel drained, I feel tired most of the time, I usually feel rather lethargic, I often feel exhausted, and I feel weary much of the time) designed to measure general feelings of tiredness and lack of energy while the remaining five items (I generally feel I have plenty of energy, I generally feel quite active, I generally feel full of vigor, I generally feel alert, and I usually feel lively) were positively worded to measure general feelings of vigor and energy (the opposite of fatigue). A single total score was computed by summing the ratings from the 10 items after reverse-coding the five positively oriented items. A higher score indicates greater feelings of chronic fatigue. In addition, separate scores for the positively and negatively worded questions were computed. The chronic fatigue questionnaire was introduced to the BCOPS study 9 months after the start of the first clinic examination and hence only 316 of the 464 participants had the opportunity to complete the questionnaire (the remaining 148 officers who did not complete the fatigue questionnaire were excluded from analyses). The instrument has high reliability with Cronbach’s alpha coefficient of 0.84 (Cohen, Manion, & Morrison, 2000). For our sample of officers the estimated alpha coefficient was 0.94 and it was obtained using the SAS procedure PROC CORR with the ALPHA option.

2.4. Assessment of on-duty injury

The second source of data (the work history records) was a longitudinal dataset that was made available in an electronic format and contained a day-by-day account of activities, for each officer, including the start time of work, the type of activity (e.g., regular work, overtime work), the type of leave (e.g., injury, sick, or vacation), and the number of hours worked for a period spanning 15 years (from May 23, 1994 to date of the BCOPS study exam). The work history records during the 1-year period prior to date of clinic examination were used to derive occurrence of injury (yes/no) for each officer and this binary variable served as the outcome variable of interest in the current analyses. For example, during the BCOPS study if an officer was examined on 8/15/2005 then we examined daily work history records of this officer from 8/15/2004 to 8/15/2005 (1-year period) to assess occurrence of on-duty injury. The work history data contained work absences due to injury that occurred while on duty. The occurrence of on-duty injury was identified when the payroll record indicates that an officer is paid for regular work but is off-duty due to injury that occurred while at work. No additional information was available concerning the type of injury or its

severity. The work history data were also used to derive dominant shift (a covariate of interest) during the same 1-year period for each officer. The methodology for derivation of dominant shift as day, afternoon or night is described in Fekedulegn et al. (2013).

2.5. Assessment of covariates

Study participants self-reported demographic and lifestyle characteristics including age, gender, race/ethnicity, marital status, education, rank, years of service, smoking and alcohol consumption, workload, physical activity, and sleep quality. Height and weight were measured with shoes removed and recorded to the nearest half centimeter and rounded up to the nearest quarter of a pound, respectively, then body mass index (BMI) was calculated as weight in kilograms divided by height in meters squared. Alcohol consumption was measured from data collected using Food Frequency Questionnaire (FFQ) where, among other things, the officers also reported how often they drank the following amounts of alcoholic beverages: beer (12 oz), red wine (6 oz), white or rose wine (6 oz), and liquor and mixed drinks (1.5 oz). The number of drinks per week was derived as the sum of consumption of these amounts from the four types of alcoholic beverages. Physical activity was assessed using the Seven-Day Physical Activity Recall questionnaire developed in the Stanford Five-City Project (Sallis et al., 1985). This was an interviewer administered questionnaire where the officers were asked to provide the number of hours they spent on three types of physical activity (occupational, sports, and household) during the previous 7 days at each of the following intensities: moderate, hard, and very hard. Sleep quality was assessed using the Pittsburgh Sleep Quality Index (PSQI) questionnaire. This was a 19-item questionnaire designed to assess the quality and pattern of sleep in adults during the previous month. A total global score for sleep quality was calculated by summing scores on the following seven components: subjective sleep quality, sleep latency, sleep duration, sleep efficiency, sleep disturbances, use of sleep medications, and daytime dysfunction. A standard cut point of ≥ 5.0 and < 5.0 was used to define “poor” and “good” sleep quality respectively (Buysse, Reynolds, Monk, Berman, & Kupfer, 1989). Workload was assessed by asking the officers the question “What is the work activity level at your district?” to which they responded by selecting one of the following: high work load (very busy with frequent complaints, high crime area); moderate work load (moderate complaint rate, average crime); or low work load (precinct not busy, low crime area).

2.6. Statistical analyses

Of the 464 BCOPS study participants, officers who did not complete the chronic fatigue questionnaire ($n = 148$) were excluded. Of the 316 remaining officers, we further excluded 36 officers who retired at least a year prior to date of exam and hence did not have work history records to determine occurrence of injury during the 1-year prior to examination. The current analyses were therefore performed using the 280 officers with complete data on both the exposure variable (chronic fatigue score) and the outcome (occurrence of on-duty injury). Initial analyses included descriptive results to characterize the composition of the study sample and examined the association of demographic and lifestyle characteristics with fatigue score and occurrence of injury using chi-square tests and analysis of variance (ANOVA).

The primary research question of interest (is there an association between chronic fatigue and occurrence of on-duty injury?) was examined using Poisson regression analysis with a robust error variance (Zou, 2004). First, we examined the association between chronic fatigue scores (overall score, score from the positive items, and score from the negative items) and occurrence of on-duty injury. In these analyses, the fatigue score was treated both as a continuous variable as well as a categorical variable (by creating tertiles). Second, we examined the association between each of the 10 items of the chronic fatigue questionnaire (individually) and injury prevalence. For this analysis we classified the ratings for each item into two categories because of small sample sizes in some of the original 5 categories. Those that rated the item as 1 (not at all) or 2 (little) were classified in one group, while those that rated the item as 3 (somewhat) or 4 (much) or 5 (very much) were combined into a second group. In both analyses prevalence ratios (PR) and their 95% confidence intervals (CI) were computed as measures of association. The unadjusted, age- and multivariate-adjusted PRs were estimated. The multivariate model adjusted for the following covariates: age, gender, race/ethnicity, education, workload, physical activity, and alcohol consumption. The variables chosen as covariates in the multivariable model are based on previous findings in the literature and those that are marginally (p -value ≤ 0.08) or significantly ($p \leq 0.05$) associated with either the exposure (fatigue) or the outcome (injury). Sleep quality and shift work are proxy measures of fatigue and hence including them in the multivariable model is considered over-adjustment. The demographic and lifestyle factors were first tested for potential effect modification by including their interaction terms in a multivariable model. For all tests, statistical significance was assessed at the 5% level. All analyses were conducted using the SAS system, version 9.3.

3. Results

3.1. Demographic and lifestyle characteristics

The demographic and lifestyle characteristics of the participants are presented in Table 1. The study population consists of 73% males and the majority was white (75%), married (70%), had a rank of patrol officer (79%), and reported high workload (64%) and poor sleep quality (57%). The mean age was 40.7 years ($SD = 6.4$) and the officers were on average overweight (mean BMI = 29.4, $SD = 4.8$). The data in Table 1 also shows mean fatigue scores and prevalence of on-duty injury (in the past year) by levels of demographic and lifestyle characteristics. Mean fatigue score did not vary significantly by levels of demographic and lifestyle characteristics except for race/ethnicity, sleep quality, and physical activity levels. White officers, and those who reported poor sleep quality had a significantly higher mean fatigue score compared to their counterparts. Physical activity hours were negatively correlated with fatigue score where officers with lower physical activity reported higher fatigue score ($r = -0.19$, p -value = 0.0017, Table 1). Prevalence of injury varied significantly by levels of workload, sleep quality, shift work, and physical activity levels (Table 1). Officers who reported high workload, poor sleep quality, and worked on the night shift had higher prevalence of on-duty injury compared to their counterparts. Physical activity hours were positively associated with prevalence of injury; for one standard deviation increase in hours of physical activity the prevalence of injury increased by 19%.

3.2. Distribution of chronic fatigue items

A majority of the officers responded favorably (“somewhat” to “very much”) to the five fatigue questionnaire items designed to measure general feelings of vigor and energy (Fig. 1A). The percentage of officers who responded “somewhat” to “very much” to these items ranged from 67.1% (I generally feel full of vigor) to 90.4% (I generally feel alert). On the other hand, the response of “somewhat” to “very much” to the five items designed to measure general feelings of tiredness and lack of energy ranged from 31.1% (I feel weary much of the time) to 39.3% (I usually feel drained). The total fatigue score (sum of score from all 10 items) ranged from 10 to 50 with a mean of 24.8 (SD = 8.1) and 46% reported above this average chronic fatigue score.

We also explored the factor structure of the chronic fatigue questionnaire for our sample of police officers. Using structural equation modeling (SEM), we fit a confirmatory factor analysis (CFA) model to estimate the latent construct (chronic fatigue) using the 10 questionnaire items. We compared a one-factor model that hypothesizes all 10 items load to a single latent variable versus a two-factor model that assumes the positively worded items load to one latent variable while the negatively worded items load to a second latent construct.

Comparison of several fit indices indicated that the two-factor model appears to fit the data better compared to the one-factor model. The Comparative Fit Index (CFI) values, where a good fit is indicated by a value of 0.95 and above, were 0.98 for the two-factor model versus 0.83 for the one-factor model. The standardized root mean square residual (SRMSR) values, where a good fit is indicated by a value lower than 0.08, were 0.024 for two-factor model and 0.082 for one-factor model.

3.3. On-duty injury

In the current sample, prevalence of on-duty injury during the past year was 24% (95% CI: 19.4–29.5); 67 of the 280 officers had on-duty injury during the past year. Among those injured, 46% (95% CI: 35.8–59.9) experienced an extended injury, which was defined as work absences that lasted at least 90 days during the past year and the average duration (in days) of injury leave was 100.6 days (SD = 100.2) (data not shown).

3.4. Association of chronic fatigue score with on-duty injury

The associations between fatigue score and occurrence of on-duty injury are presented in Table 2. Results indicate an increasing trend in prevalence of on-duty injury across tertiles of total fatigue score (19.6%, 21.7%, 30.8% for the lowest, middle, and highest tertiles, respectively, trend p -value = 0.0372). After adjusting for age, gender, race/ethnicity, education, workload, physical activity, and alcohol consumption, the prevalence of injury among officers in the highest tertile of total fatigue score was 67% larger compared to those in the lowest tertile (PR = 1.67, 95% CI: 0.99–2.83, p = 0.0554) but with borderline statistical significance. Analyses of total fatigue score in continuous form indicated that a 5-unit increase in total fatigue score was associated with a 12% increase in prevalence of on-duty injury (PR = 1.12, 95% CI: 0.99–1.27, p = 0.0746) yet the estimate had borderline statistical significance.

The association between fatigue score from the positively worded items and on-duty injury achieved statistical significance. A 5-unit increase in fatigue score of the positively worded items was associated with a 33% increase in prevalence of injury (PR = 1.33, 95% CI: 1.04–1.70, $p = 0.0215$). Injury prevalence was 62% higher among officers in the highest tertile of the score for the positive items compared to those in the lowest tertile (PR = 1.62, 95% CI: 0.94–2.81, $p = 0.0827$). On the other hand, there was no significant association between fatigue score from the negatively worded items and occurrence of injury; a 5-unit increase in score from these items was associated with 13% increase in prevalence of injury (PR = 1.13, 95% CI: 0.90–1.41, $p = 0.2842$), and those in the highest tertile of fatigue score from the negative items had injury prevalence that was 45% larger compared to those in the lowest tertile (PR = 1.45, 95% CI: 0.83–2.53, $p = 0.1941$).

3.5. Association of individual items of chronic fatigue questionnaire with on-duty injury

The association of ratings to the individual items of the chronic fatigue questionnaire with occurrence of injury is presented in Table 3 (for the positively worded items) and Table 4 (for the negatively worded items). As expected, the results in Table 3 show a significantly higher prevalence of on-duty injury among officers who responded “not at all” or “little” to 4 of the 5 positively worded items. After multivariate adjustment, officers who do not feel active (PR = 1.74, 95% CI: 1.09–2.80, $p = 0.0210$) had significantly higher prevalence of on-duty injury compared to their counterparts. Prevalence of on-duty injury was 75% larger among officers who generally do not feel full of vigor compared to those who do feel full of vigor (PR = 1.75, 95% CI: 1.13–2.71, $p = 0.0118$). Officers who generally do not feel alert had more than double the prevalence of injury compared to those who do feel alert (PR = 2.31, 95% CI: 1.36–3.94, $p = 0.0020$). Officers who do not feel lively had injury prevalence that is 67% larger (PR = 1.67, 95% CI: 1.00–2.79, $p = 0.0495$) compared to their counterparts. On the other hand, there was no significant association between the ratings for the negatively worded items and prevalence of injury (Table 4); the only exception is a borderline statistical significance between the rating for “I often feel exhausted” and injury occurrence where those who often felt exhausted had a 52% higher prevalence of injury compared to their counterparts (PR = 1.52, 95% CI: 0.99–2.33, $p = 0.0541$).

4. Discussion

There are approximately 900,000 sworn law enforcement officers serving in the United States (NLEOMF, 2016). This workforce is known to disproportionately suffer from cardiovascular, metabolic, and psychosocial disorders (Barron, 2008; Hartley, Burchfiel, Fekedulegn, Andrew, & Violanti, 2011; Liberman et al., 2002; Violanti et al., 2009; Zimmerman, 2012). In addition, policing is one of the occupations with the highest rate of non-fatal on-duty injury. According to the Bureau of Labor Statistics (BLS), police and sheriff's patrol officers was one of six occupations (correctional officers and jailers, firefighters, nursing assistants, construction laborers, and heavy and tractor-trailer truck drivers) where the incidence rate of non-fatal workplace injury, per 10,000 full-time workers, was greater than 300 (BLS, 2014). The rate of non-fatal occupational injuries is two to three times the national average, while fatal injury rates are nearly four times greater among officers compared to the average American worker (LaTourrette, 2011). Although

fatigue is commonly understood to be a risk factor for on-duty injury among police, scientific literature is limited. Assessment of the prevalence of fatigue among law enforcement officers, who work under high-risk and dynamic environments, engage in extended driving, and often need to make on-the-spot decisions in complex and ambiguous situations, is particularly important because fatigue in police officers can have devastating consequences to the officers and the general public (Vila & Kenney, 2002). Police officers, therefore, represent a unique occupation group for research focusing on fatigue, sleep, and human performance (Vila, 2006). In this study, we assessed chronic fatigue in police officers working in a mid-sized urban department and examined its association with occurrence of non-fatal workplace injury. The results indicated that 46% of the officers had above average chronic fatigue score (N24.8) and nearly 40% reported feeling drained. Overall, the prevalence of workplace injury increased significantly across tertiles of total fatigue score. In particular, injury was more than twice as prevalent among officers who generally did not feel alert compared to their counterparts. Prevalence of non-fatal workplace injury was at least 65% larger among officers who did not feel active, full of vigor, or lively compared to those who did.

There are a number of studies that attempted to examine the relationship between fatigue and safety outcomes in working populations. Some were based on assessment of fatigue using validated questionnaires (Fang, Jiang, Zhang, & Wang, 2015; Swaen, Van Amelsvoort, Bültmann, & Kant, 2003) while most use proxy indicators of fatigue (Kao, Spitzmueller, Cigularov, & Wu, 2016; Salminen et al., 2010) namely sleep deprivation/disorder, shift work, and irregular work hours. In a prospective study (Swaen et al., 2003) that examined the effect of fatigue [assessed using the Checklist Individual Strength (CIS) questionnaire] on injury, the risk of workplace injury was 69% higher among those in the highest tertile of fatigue score compared to those in the lowest tertile following adjustment for multiple potential confounders (RR = 1.69, 95% CI: 1.03–2.78). The magnitude of the effect size is consistent with our findings despite differences in study design and study population. The study by Swaen et al. (2003) also reported the negative consequence of irregular work hours; those working irregular shifts (but not night shift) had a five-fold higher risk (crude RR = 4.76, 95% CI: 2.42–9.35) of occupational injuries compared to those working on day shifts. There are numerous studies that showed sleep problems (another proxy measure of fatigue) are associated with increased risks of workplace injuries and accidents (Chau, Mur, Tournon, Benamghar, & Dehaene, 2004; Salminen et al., 2010; Uehli et al., 2014a, 2014b). A recent meta-analysis of observational studies dealing with sleep problems and injury revealed that workers with sleep problems were 62% more likely to being injured at the workplace compared to those without sleep problem (RR = 1.62, 95% CI: 1.43–1.84) and that 13% of the work-place injuries could be attributed to sleep problems (Uehli et al., 2014a). A case-control study (Chau et al., 2004) that examined correlates of occupational injury indicated that injured workers had 30% higher odds of having a sleep disorder compared to non-injured workers (OR = 1.30, 95% CI: 1.08–1.57). A prospective study of Finish public sector workers (Salminen et al., 2010) also reported a similar estimate where the risk of workplace injury was 38% higher among those who experienced disturbed sleep compared to those who did not (OR = 1.38, 95% CI: 1.02–1.87). Among truck drivers long workhours (driving and non-driving) have been significantly associated with an increase in safety critical events

(Socolich et al., 2013), and driver fatigue was a significant risk factor for occupational light vehicle crashes (OR = 2.1, 95% CI: 1.5–2.7; Stuckey, Glass, LaMontagne, Wolfe, & Sim, 2010).

In policing, formal studies of fatigue and on-duty injury, although limited and based on proxy indicators of fatigue, highlight findings consistent with those in other occupational groups or the general population. An experimental study by James and Vila (2015) examined the extent fatigue (defined as shift work) degrades post-shift non-operational driving (i.e., lab-based simulated driving) performance of officers. Their results showed that officers working on night shift (fatigued condition) had significantly greater lane deviation compared to those on day shift. Analysis of shift work in relation to injury based on the same study population used for the current study (Violanti et al., 2012) showed that the incidence rate of non-fatal on-duty injury was 72% larger for officers working on the night shift compared to those on the day shift (IRR = 1.72; 95% CI = 1.26–2.36) while the risk of long-term injury (90 days of work absence) was three-fold higher among officers working the night shift (Violanti et al., 2013). These studies also indicated that high workload in combination with night shift work significantly exacerbated the risk of workplace injury. Officers in the United States are reported to work extraordinarily high numbers of hours per week (graveyard shifts, overtime, second jobs, etc.) resulting in insufficient sleep and poor rest that heightens the risk of injury (Senjo, 2011). The prevalence of poor sleep quality in U.S. police officers was 64% compared to 45% among those not involved in emergency services (Neylan et al., 2002). In a study of police officers from the United States and Canada, Rajaratnam et al. (2011) reported that 40% had at least one sleep disorder, 34% had obstructive sleep apnea, 7% had insomnia, 29% reported excessive sleepiness, and 26% reported falling asleep while driving at least once in a month. In policing, where officers are engaged in extended driving, nodding off while driving and difficulty to maintain constant vigilance because of sleepiness leads to disastrous outcomes to both the officers and the public at large. A study by Vila (2000) reported that 4 of out of 8 officers involved in on-duty injuries and accidents were impaired because of fatigue. The most common types of nonfatal injuries among officers are strains and sprains, particularly those to the back, often caused by trips and falls (particularly during foot pursuits) and extended driving. Vehicle crashes represent the greatest fraction of both fatal and non-fatal injuries, making driving the most dangerous activity police engage in (LaTourrette, 2011).

Fatigue has also been reported to be associated with (or co-occurs with) a number of adverse psychological health outcomes including depression and anxiety, and chronic diseases (Chen, 1986; Franssen, Bültmann, Kant, & Van Amelsvoort, 2003). U.S. workers reporting fatigue are four times more likely to experience depressive symptoms than workers who did not report fatigue (Ricci et al., 2007). In our police study sample, a 5-unit increase in total fatigue score was associated with a 72% elevation in prevalence of depression (PR = 1.72, 1.49–1.98, $p < 0.0001$, data not shown). Overall, the proportion of officers who self-reported their general health as “fair or poor” increased by 51% for a 5-unit increase in total fatigue score (PR = 1.51, 1.24–1.84, $p < 0.0001$, data not shown). It is worth mentioning that worker fatigue also has significant economic consequences. The health related lost productive time (LPT) among workers with fatigue cost U.S. employers an estimated \$136 billion annually

(Ricci et al., 2007); where 66% of U.S. workers with fatigue reported health-related lost productive time (LPT) compared to only 26% of those without fatigue.

The current study has several strengths including the use of objective daily work history records from which on-duty injury was ascertained, a relatively large sample, high reliability of the chronic fatigue questionnaire which was specifically designed to assess fatigue in occupations involving shift work (Khaleghipour, Masjedi, & Kelishadi, 2015), and adjustment of the association of interest for multiple potential confounders. Despite these strengths, the findings from this study ought to be interpreted in the context of potential limitations. The study is based on urban police officers from the eastern United States (convenience sample) and therefore may have limited generalizability to all officers. Chronic fatigue was assessed through self-report and hence there is a possibility of response bias (especially socially desirable responding) that could underestimate the prevalence of fatigue. Despite our effort, we were not able to access the data on the type and severity of injury for reasons related to privacy. From a methodological viewpoint, occurrence of on-duty injury was assessed during the 1-year prior to assessment of fatigue, and therefore based on the cross-sectional study design it is assumed that chronic fatigue assessed at the clinic examination has been present or consistent throughout the previous 1 year period. The cross-sectional study design limits casual inference.

In summary, this study of urban police officers showed that those who do not feel active, full of vigor, alert, or lively had a significantly higher prevalence of non-fatal work place injury compared to their counter parts. To the authors' best knowledge, there are no epidemiologic studies that assessed chronic fatigue using one of available validated instruments and examined its association with objectively assessed on-duty injury among police officers in the United States. This study adds to the body of knowledge regarding the association of fatigue with workplace injury among high stress occupations and may have future implications that ultimately will lead to interventions that could reduce officer's fatigue and injury occurrence. Policing, by its nature, involves exposure to inherent physical and psychological dangers as well as additional occupational stressors (e.g., long work hours, shift work, and irregular schedules) that significantly increase the risk for chronic fatigue and numerous adverse health outcomes including fatal and non-fatal workplace injuries. In their report titled "Tired cops" Vila and Kenney (2002) provide accounts of devastating tragedies (injuries) associated with police fatigue. Altogether, the human and economic cost associated with fatigue and workplace injury could represent a substantial burden to officers and their families and warrants greater attention. Hence, there is a need to identify strategies to reduce the risk factors that lead to chronic fatigue and workplace injury. Comprehensive fatigue management programs that include education on the health and safety consequences of fatigue, regulations on the length of work hours per day and per week, workplace interventions that improve alertness/fitness, and screening for sleep disorders are essential to minimize fatigue and its negative consequences. Fatigue intervention also ought to consider psychosocial work characteristics. For example, a prospective study has shown that decision latitude in men and co-worker social support in women were protective against fatigue (Bültmann, Kant, Van Den Brandt, & Kasl, 2002). Physical inactivity and obesity have long been recognized as risk factors for fatigue (Chen, 1986). A study by Zimmerman (2012) indicated that workplace programs to promote the health and fitness of police officers are

commonly lacking despite the fact that obesity may be more common in police officers compared with other groups. Weight loss and fitness could be important factors in reducing acute musculoskeletal injuries (LaTourrette, 2011). Future studies with larger sample size and a prospective design are worthwhile and could provide better insight in designing effective interventions.

Acknowledgments

This work was supported by the National Institute for Occupational Safety and Health (NIOSH), contract no. 200-2003-01580. The findings and conclusions in this article are those of the authors and do not necessarily represent the views of the National Institute for Occupational Safety and Health. In addition, citations to websites external to NIOSH do not constitute NIOSH endorsement of the sponsoring organizations or their programs or products. Furthermore, NIOSH is not responsible for the content of these websites. All web addresses referenced in this document were accessible as of the publication date. This project was made possible through a partnership with the CDC Foundation.

Biographies

Fekedulegn Desta PhD, is a statistician at the Biostatistics and Epidemiology Branch in the Health Effects Laboratory Division of the National Institute for Occupational Safety and Health at CDC. He received a MS in statistics and a Ph.D. in forest science from West Virginia University. He served as a research assistant professor at West Virginia University department of statistics for several years before joining NIOSH. His research interests include analytic methods in epidemiology, actigraphy based assessment of sleep and physical activity, and impact of occupational stress on health.

Cecil M. Burchfiel received his MPH and PhD degrees in epidemiology at the University of Michigan School of Public Health. He has worked as an epidemiologist with the Veterans Administration, the Department of Preventive Medicine and Biometrics at the University of Colorado School of Medicine, the National Heart, Lung and Blood Institute of NIH, and currently is Chief of the Biostatistics and Epidemiology Branch in the Health Effects Laboratory Division of the National Institute for Occupational Safety and Health at CDC. His research interests focus on cardiovascular disease epidemiology with emphasis on associations of workplace stressors with subclinical cardiovascular and metabolic disorders.

Claudia C. Ma earned her MS degree in Agronomy in P.R. China and worked more than ten years with the Chinese government in forest research and landscape project management. After receiving her MPH in the United States, she has worked as an Epidemiologist seven years in the Biostatistics and Epidemiology Branch, Health Effects Laboratory Division at NIOSH. Her research focuses on occupational exposures and the risks of cardio-metabolic outcomes, musculoskeletal disorders, and the potential role of parental occupational exposures in autism in working population.

Michael E Andrew PhD, serves as senior statistician for the Biostatistics and Epidemiology Branch of the Health Effects Laboratory Division, of the National Institute for Occupational Safety and Health (NIOSH) at the Centers for Disease Control and Prevention (CDC). His research interests include cardiovascular disease epidemiology with recent focus on

associations of workplace stressors with autonomic dysfunction as measured by heart rate variability, and protective factors related to workplace stressors.

Tara A. Hartley PhD, MPA, MPH is an Epidemiologist in the Biostatistics and Epidemiology Branch, Health Effects Laboratory Division of the National Institute for Occupational Safety and Health. Dr. Hartley earned masters' degrees in public administration and public health and a doctoral degree in public health sciences from West Virginia University. She has been a member of the BCOPS Study research team since 2004. Her research interests have focused on the role of workplace stress in the development of subclinical cardiovascular disease, and the use of novel biomarkers for identifying cardiovascular and metabolic diseases in epidemiologic research.

Luenda E. Charles PhD, MPH is a senior Epidemiologist/Health Scientist with the National Institute for Occupational Safety and Health at the Centers for Disease Control and Prevention. She obtained a Master of Public Health degree in Occupational and Environmental Health from Emory University, Atlanta, Georgia, and a PhD in Epidemiology from the University of North Carolina, Chapel Hill, North Carolina. Previous research involved occupational and environmental factors associated with cancer. Her current research focuses on investigations of occupational exposures with sub-clinical cardiovascular disease and other health problems.

Ja K. Gu MSPH, is a statistician in the Biostatistics and Epidemiology Branch, Health Effects Laboratory Division, National Institute for Occupational Safety and Health. He received a MSPH in Biostatistics from the University of South Carolina at Columbia and worked traumatic brain injury and spinal cord injury cohort research in South Carolina Department of Disability and Special Needs and Medical University of South Carolina for 8 years. He has worked with the Buffalo Cardio-Metabolic Occupational Police Stress (BCOPS) Study since 2008, providing data management assistance and statistical analysis. In addition, he has provided guidance and assistance to NIOSH scientists in statistical methods. His research interests are obesity, injury, and physical activity in US workers, especially workers in the protective service.

John M. Violanti PhD, is a research professor in the department of Epidemiology & Environmental Health, State University of NY at Buffalo, NY. His research focuses primarily on police stress and trauma, health and suicide.

References

- Barron S (2008). Occupational stress: The emerging threat to police officers. *Journal of Occupational Health and Safety - Australia and New Zealand*, 24(6), 553–561.
- Barton J, Spelten E, Totterdell P, Smith L, Folkard S, & Costa G (1995). The Standard Shiftwork Index: A battery of questionnaires for assessing shiftwork-related problems. *Work and Stress*, 9(1), 4–30.
- Berecki-Gisolf J, Smith PM, Collie A, & McClure RJ (2015). Gender differences in occupational injury incidence. *American Journal of Industrial Medicine*, 58(3), 299–307. [PubMed: 25641425]
- Bültmann U, Kant IJ, Van Den Brandt PA, & Kasl SV (2002). Psychosocial work characteristics as risk factors for the onset of fatigue and psychological distress: respective results from the Maastricht cohort study. *Psychological Medicine*, 32(2), 333–345. [PubMed: 11871373]

- Bureau of Labor Statistics [BLS] (2014). Nonfatal occupational injuries and illnesses involving days away from work by selected worker occupation and ownership, 2014. (Table.4 Number, incidence rate, and median days away from work). (<http://www.bls.gov/news.release/osh2.t04.htm> Last accessed 10 February 2016).
- Buysse DJ, Reynolds CF, 3rd, Monk TH, Berman SR, & Kupfer DJ (1989). The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Research*, 28, 193–213. [PubMed: 2748771]
- Caban-Martinez AJ, Courtney TK, Chang W, Lombardi DA, Huang Y, Brennan MJ, ... Verma SK (2015). Leisure-time physical activity, falls, and fall injuries in middle-aged adults. *American Journal of Preventive Medicine*, 49(6), 888–901. [PubMed: 26232899]
- Chau N, Mur J, Tournon C, Benamghar L, & Dehaene D (2004). Correlates of occupational injuries for various jobs in railway workers: A case-control study. *Journal of Occupational Health*, 46(4), 272–280. [PubMed: 15308826]
- Chen MK (1986). The epidemiology of self-perceived fatigue among adults. *Preventive Medicine*, 15(1), 74–81. [PubMed: 3714661]
- Cohen L, Manion L, & Morrison K (2000). *Research methods in education* (5th ed.). London: Routledge, 211.
- De Vries J, Michielsen HJ, & Van Heck GL (2003). Assessment of fatigue among working people: A comparison of six questionnaires. *Occupational and Environmental Medicine*, 60(Suppl. 1), i10–i15. [PubMed: 12782741]
- Fang D, Jiang Z, Zhang M, & Wang H (2015). An experimental method to study the effect of fatigue on construction workers' safety performance. *Safety Science*, 73, 80–91.
- Fekedulegn D, Burchfiel CM, Hartley TA, Andrew ME, Charles LE, Tinney-Zara CA, & Violanti JM (2013). Shiftwork and sickness absence among police officers: The BCOPS study. *Chronobiology International*, 30(7), 930–941. [PubMed: 23808812]
- Franssen PML, Bültmann U, Kant I, & Van Amelsvoort LGPM (2003). The association between chronic diseases and fatigue in the working population. *Journal of Psychosomatic Research*, 54(4), 339–344. [PubMed: 12670611]
- Garbarino S, Repice AM, Traversa F, Spigno F, Mascialino B, Mantineo G, ... Bonsignore AD (2007). Commuting accidents: The influence of excessive daytime sleepiness. A review of an Italian police officers population. [Infortuni in itinere: Ruolo della sonnolenza in una popolazione di operatori della polizia di stato]. *Giornale Italiano di Medicina del Lavoro ed Ergonomia*, 29(3), 324–326. [PubMed: 18409708]
- Hartley TA, Burchfiel CM, Fekedulegn D, Andrew ME, & Violanti JM (2011). Health disparities in police officers: Comparisons to the U.S. general population. *International Journal of Emergency Mental Health*, 13(4), 211–220. [PubMed: 22900455]
- Hurley DT, & Lebbon AR (2012). A comparison of nonfatal occupational injuries and illnesses among Hispanic versus non-Hispanic workers in the United States. *Hispanic Journal of Behavioral Sciences*, 34(3), 474–490.
- James SM, & Vila B (2015). Police drowsy driving: Predicting fatigue-related performance decay. *Policing*, 38(3), 517–538.
- Kao K, Spitzmueller C, Cigularov K, & Wu H (2016). Linking insomnia to workplace injuries: A moderated mediation model of supervisor safety priority and safety behavior. *Journal of Occupational Health Psychology*, 21(1), 91–104. [PubMed: 26011243]
- Khaleghipour S, Masjedi M, & Kelishadi R (2015). Circadian type, chronic fatigue, and serum IgM in the shift workers of an industrial organization. *Advances in Biology Research*, 4(61).
- Kim SK, Kim H, Lee K, Kang HT, Oh SS, & Ko SB (2014). The relationship between injury and socioeconomic status in reference to the Fourth Korean National Health and Nutrition Examination Survey. *Annals of Occupational and Environmental Medicine*, 26(1), 1. [PubMed: 24472308]
- Landen DD, & Hendricks SA (1992). Estimates from the national health interview survey on occupational injury among older workers in the United States. *Scandinavian Journal of Work, Environment and Health*, 18(Suppl. 2), 18–20.
- LaTourrette T (2011). Safety and health protection efforts in the police service. *The Police Chief*, 78, 74–78.

- Lewis G, & Wessely S (1992). The epidemiology of fatigue: More questions than answers. *Journal of Epidemiology and Community Health*, 46(2), 92–97. [PubMed: 1583440]
- Lieberman AM, Best SR, Metzler TJ, Fagan JA, Weiss DS, & Marmar CR (2002). Routine occupational stress and psychological distress in police. *Policing*, 25(2), 421–441.
- Nakata A, Ikeda T, Takahashi M, Haratani T, Hojou M, Fujioka Y, ... Araki S (2006). Impact of psychosocial job stress on non-fatal occupational injuries in small and medium sized manufacturing enterprises. *American Journal of Industrial Medicine*, 49(8), 658–669. [PubMed: 16758484]
- National Law Enforcement Officers Memorial Fund [NLEOMF] (2016). Law enforcement facts. <http://www.nleomf.org/facts/enforcement/> (Last accessed 10 February 2016).
- National Occupational Research Agenda [NORA] (2009). National public safety agenda for occupational safety and health research and practice in the U.S public safety sub sector. <http://www.cdc.gov/niosh/nora/comment/agendas/pubsafsub/pdfs/PubSafSubApr2009.pdf> (Last accessed 19 February 2016).
- Neylan TC, Metzler TJ, Best SR, Weiss DS, Fagan JA, Liberman A, ... Marmar CR (2002). Critical incident exposure and sleep quality in police officers. *Psychosomatic Medicine*, 64(2), 345–352. [PubMed: 11914452]
- Rajaratnam SMW, Barger LK, Lockley SW, Shea SA, Wang W, Landrigan CP, ... Czeisler CA (2011). Sleep disorders, health, and safety in police officers. *JAMA : The Journal of the American Medical Association*, 306(23), 2567–2578. [PubMed: 22187276]
- Ricci JA, Chee E, Lorandean AL, & Berger J (2007). Fatigue in the U.S. workforce: Prevalence and implications for lost productive work time. *Journal of Occupational and Environmental Medicine*, 49(1), 1–10. [PubMed: 17215708]
- Sallis JF, Haskell WL, Wood PD, Fortmann SP, Rogers T, Blair SN, & Paffenbarger RS, Jr. (1985). Physical activity assessment methodology in the Five-City Project. *American Journal of Epidemiology*, 121, 91–106. [PubMed: 3964995]
- Salminen S, Oksanen T, Vahtera J, Sallinen M, Härmä M, Salo P, ... Kivimäki M (2010). Sleep disturbances as a predictor of occupational injuries among public sector workers. *Journal of Sleep Research*, 19(1 PART. 2), 207–213. [PubMed: 19840241]
- Senjo SR (2011). Dangerous fatigue conditions: A study of police work and law enforcement administration. *Police Practice and Research*, 12(3), 235–252.
- Soccolich SA, Blanco M, Hanowski RJ, Olson RL, Morgan JF, Guo F, & Wu S (2013). An analysis of driving and working hour on commercial motor vehicle driver safety using naturalistic data collection. *Accident Analysis and Prevention*, 58, 249–258. [PubMed: 22818778]
- Stallones L, & Xiang H (2003). Alcohol consumption patterns and work-related injuries among Colorado farm residents. *American Journal of Preventive Medicine*, 25(1), 25–30. [PubMed: 12818306]
- Stuckey R, Glass DC, LaMontagne AD, Wolfe R, & Sim MR (2010). Risk factors for worker injury and death from occupational light vehicles crashes in New South Wales (Australia). *American Journal of Industrial Medicine*, 53(9), 931–939. [PubMed: 20698024]
- Swaen GMH, Van Amelsvoort LGPM, Bültmann U, & Kant IJ (2003). Fatigue as a risk factor for being injured in an occupational accident: Results from the Maastricht cohort study. *Occupational and Environmental Medicine*, 60(Suppl. 1), i88–i92. [PubMed: 12782753]
- Uehli K, Mehta AJ, Miedinger D, Hug K, Schindler C, Holsboer-Trachsler E, ... Künzli N (2014a). Sleep problems and work injuries: A systematic review and meta-analysis. *Sleep Medicine Reviews*, 18, 61–73. [PubMed: 23702220]
- Uehli K, Miedinger D, Bingisser R, Dürr S, Holsboer-Trachsler E, Maier S, ... Leuppi JD (2014b). Sleep quality and the risk of work injury: A Swiss case-control study. *Journal of Sleep Research*, 23(5), 545–553. [PubMed: 24889190]
- Vila BJ (2000). Tired cops: the importance of managing police fatigue. Washington DC: Police Executive Research Forum.
- Vila B (2006). Impact of long work hours on police officers and the communities they serve. *American Journal of Industrial Medicine*, 49(11), 972–980. [PubMed: 17006951]

- Vila B, & Kenney DJ (2002). Tired cops: The prevalence and potential consequences of police fatigue. *NIJ Journal*, 248, 16–21.
- Violanti JM, Burchfiel CM, Hartley TA, Mnatsakanova A, Fekedulegn D, Andrew ME, ... Vila BJ (2009). Atypical work hours and metabolic syndrome among police officers. *Archives of Environmental and Occupational Health*, 64(3), 194–201. [PubMed: 19864222]
- Violanti JM, Burchfiel CM, Miller DB, Andrew ME, Dorn J, Wactawski-Wende J, ... Trevisan M (2006). The buffalo cardio-metabolic occupational police stress (BCOPS) pilot study: Methods and participant characteristics. *Annals of Epidemiology*, 16(2), 148–156. [PubMed: 16165369]
- Violanti JM, Fekedulegn D, Andrew ME, Charles LE, Hartley TA, Vila B, & Burchfiel CM (2012). Shift work and the incidence of injury among police officers. *American Journal of Industrial Medicine*, 55(3), 217–227. [PubMed: 22228219]
- Violanti JM, Fekedulegn D, Andrew ME, Charles LE, Hartley TA, Vila B, & Burchfiel CM (2013). Shift work and long-term injury among police officers. *Scandinavian Journal of Work, Environment and Health*, 39(4), 361–368.
- Waggoner LB, Grant DA, Van Dongen HPA, Belenky G, & Vila B (2012). A combined field and laboratory design for assessing the impact of night shift work on police officer operational performance. *Sleep*, 35(11), 1575–1577. [PubMed: 23115407]
- Zimmerman FH (2012). Cardiovascular disease and risk factors in law enforcement personnel: A comprehensive review. *Cardiology in Review*, 20(4), 159–166. [PubMed: 22314143]
- Zou G (2004). A modified poisson regression approach to prospective studies with binary data. *American Journal of Epidemiology*, 159(7), 702–706. [PubMed: 15033648]




Fig. 1. Percentage of participants responding to chronic fatigue questionnaire items among police officers. Responses of “somewhat” to “very much” were combined. Part A shows percentages for the positively worded items (I generally feel I have plenty of energy, I generally feel quite active, I generally feel full of vigor, I generally feel alert, and I usually feel lively). Part B shows percentages for the negatively worded items (I usually feel drained, I feel tired most of the time, I usually feel rather lethargic, I often feel exhausted, and I feel weary much of the time).

Demographic and life style characteristics and their association with fatigue score and on-duty injury, BCOPS study, 2004–2009.

Table 1

Characteristics	Study sample		Chronic fatigue score		On-duty injury (past year)	
	N	% (±SD)	Mean ± SD	p-Value ^c	Prevalence (%)	p-Value ^d
<i>Gender</i>						
Male	205	73.2	24.2 ± 7.3	0.0598	24.4	0.7647
Female	75	26.8	26.3 ± 9.9		22.7	
<i>Race</i>						
White	205	74.6	25.6 ± 8.5	0.0029	23.4	0.6973
Black/Hispanic	70	25.5	22.2 ± 6.6		25.7	
<i>Education</i>						
High school/GED	27	9.7	21.7 ± 7.6	0.0753	11.1	0.2664
College <4 yrs	158	56.8	25.4 ± 8.6		25.3	
College 4+ yrs	93	33.5	24.4 ± 7.3		24.7	
<i>Marital status</i>						
Single	36	13.0	25.1 ± 8.3	0.7917	33.3	0.1608
Married	194	59.8	24.9 ± 8.2		20.6	
Divorced	48	17.3	24.0 ± 8.0		29.2	
<i>Smoking status</i>						
Current	53	19.2	26.6 ± 9.6	0.1172	26.4	0.3460
Former	50	18.1	23.3 ± 8.0		16.0	
Never	173	52.7	24.7 ± 7.7		25.4	
<i>Rank</i>						
Patrol officer	220	78.9	26.0 ± 8.0	0.1938	24.6	0.4996
Other ^a	59	21.2	24.4 ± 8.2		20.3	
<i>Workload (high)</i>						
Low/medium	98	35.8	25.3 ± 8.5	0.4273	16.3	0.0317
High	176	64.2	24.5 ± 8.0		27.8	
<i>Sleep quality</i>						
Poor	152	57.4	27.1 ± 8.1	<0.0001	28.3	0.0452
Good	113	42.6	21.6 ± 7.3		17.7	

Characteristics	Study sample		Chronic fatigue score		On-duty injury (past year)	
	N	% (±SD)	Mean ± SD	p-Value ^c	Prevalence (%)	p-Value ^d
<i>Shift work (past year)</i>						
Day	117	42.2	24.8 ± 9.1	0.9451	18.8	0.0019
Afternoon	86	31.1	25.0 ± 7.5		16.3	
Night	74	26.7	24.5 ± 7.4		37.8	
Age (in years)	280	40.7 ± 6.4	r ^e = -0.09	0.1340	PR ^f = 0.87	0.2071
Years of service	279	13.7 ± 6.5	r = -0.11	0.0696	PR = 0.78	0.0531
Body mass index (kg/m ²)	279	29.4 ± 4.8	r = 0.08	0.1574	PR = 0.89	0.2782
Physical activity ^b (h/week)	279	16.2 ± 14.3	r = -0.19	0.0017	PR = 1.19	0.0489
No. of alcohol drinks/week	275	5.2 ± 8.4	r = 0.10	0.0909	PR = 1.02	0.8934

Results for continuous variables are means ± SD.

^aOther includes Sergeant, Lieutenant, Captain, and Detective.

^bPhysical activity hours per week including occupational, household, and leisure time activities.

^cp-Values are from analysis of variance (ANOVA) for categorical variables or correlation analysis for continuous variables.

^dp-Values are from χ^2 tests of independence or Fisher's exact test for categorical variables. For continuous variables, P-values are from Poisson regression testing linear trend in prevalence of injury.

^eDenotes the correlation between fatigue score and each of the continuous variables.

^fPR denotes the prevalence ratio associated with one standard deviation increase in each continuous variable.

Table 2

Prevalence and prevalence ratio (95% CI) of on-duty injury by levels of chronic fatigue score (overall score, score for the positive items, and score for the negative items).

	N	Number of injury cases	Prevalence of injury (%)	Prevalence ratio (95% CI)		
				Unadjusted	Age-adjusted	MV-adjusted ¹
Total fatigue score (all 10 items) ^b	280	67	23.9 (19.4–29.5)	1.08 (0.96–1.22)	1.07 (0.96–1.21)	1.12 (0.99–1.27)
Tertiles						
Low (10–20)	97	19	19.6(13.1–29.3)	Referent		
Medium (21–27)	92	20	21.7 (14.8–32.0)	1.11 (0.63–1.94)	1.13 (0.64–1.98)	1.08 (0.61–1.91)
High (28–50)	91	28	30.8 (22.6–41.9) (p = 0.0372) ^c	1.57 (0.95–2.61)	1.54 (0.93–2.55)	1.67 (0.99–2.83)
Score for positive items ^b				1.21 (0.96–1.52)	1.20 (0.95–1.50)	1.33 (1.04–1.70)
Tertiles						
Low (5–10)	85	17	20.0(13.1–30.6)	Referent		
Medium (11–15)	115	27	23.5 (16.9–32.7)	1.17 (0.69–2.01)	1.13 (0.66–1.94)	1.06(0.61–1.83)
High (16–25)	SO	23	28.8 (20.4–40.6) (p = 0.0945) ^c	1.44 (0.83–2.49)	1.40(0.81–2.41)	1.62 (0.94–2.81)
Score for negative items ^b				1.09 (0.88–1.35)	1.08 (0.87–1.33)	1.13 (0.90–1.41)
Tertiles						
Low (5–8)	86	17	19.8 (12.9–30.3)	Referent		
Medium (9–13)	108	27	25.0(18.0–34.7)	1.26(0.74–2.16)	1.27 (0.74–2.18)	1.35 (0.78–2.31)
High (14–25)	86	23	26.7 (18.9–37.9) (p = 0.1418) ^c	1.35 (0.78–2.35)	1.33 (0.77–2.30)	1.45 (0.83–2.53)

^a Adjusted for age, gender, race/ethnicity, education, workload, physical activity, and alcohol consumption.

^b Prevalence ratios are for 5-unit increase in fatigue score.

^c P-value testing a linear trend in prevalence of injury across increasing tertiles of chronic fatigue score (Cochran–Armitage trend test).

Table 3

Prevalence and prevalence ratio (95% CI) of on-duty injury by levels of the five positively worded items of the chronic fatigue questionnaire.

Characteristics	Number participants	Number of injury cases	Prevalence of injury (%)	Prevalence ratio (95% CI)		
				Unadjusted	Age-adjusted	MV-adjusted ^a
<i>I generally feel I have plenty of energy</i>						
Somewhat – very much	235	54	23.0	Referent		
Not at all – little	45	13	28.9	1.26 (0.75–2.10)	1.26 (0.75–2.10)	1.43 (0.85–2.40) ^b (p = 0.1732)
<i>I generally feel quite active</i>						
Somewhat – very much	227	50	22.0	Referent		
Not at all – little	53	17	32.1	1.46 (0.92–2.31)	1.42 (0.89–2.25)	1.74 (1.09–2.80) (p = 0.0210)
<i>I generally feel full of vigor</i>						
Somewhat – very much	188	38	20.2	Referent		
Not at all – little	92	29	31.5	1.56 (1.03–2.36)	1.55 (1.02–2.34)	1.75 (1.13–2.71) (p = 0.0118)
<i>I generally feel alert</i>						
Somewhat – very much	253	56	22.1	Referent		
Not at all – little	27	11	40.7	1.84 (1.11–3.07)	1.81 (1.08–3.01)	2.31 (1.36–3.94) (p = 0.0020)
<i>I usually feel lively</i>						
Somewhat – very much	231	52	22.5	Referent		
Not at all – little	49	15	30.6	1.36 (0.84–2.21)	1.33 (0.82–2.15)	1.67 (1.00–2.79) (p = 0.0495)

^a Adjusted for age, gender, race/ethnicity, education, workload, physical activity, and alcohol consumption.

^b p-Value for the multivariable adjusted prevalence ratio.

Table 4

Prevalence and prevalence ratio (95% CI) of on-duty injury by levels of the five negatively worded items of the chronic fatigue questionnaire.

Characteristics	Number participants	Number of injury cases	Prevalence of injury (%)	Prevalence ratio (95% CI)		
				Unadjusted	Age adjusted	MV-adjusted ^a
<i>I usually feel drained</i>						
Not at all – little	170	39	22.9	Referent		
Somewhat – very much	110	28	25.5	1.11 (0.73–1.69)	1.08 (0.71–1.65)	1.13 (0.74–1.73) ^b (p = 0.5667)
<i>I feel tired most of the time</i>						
Not at all – little	170	37	21.8	Referent		
Somewhat – very much	110	30	27.3	1.25 (0.83–1.90)	1.24 (0.82–1.89)	1.33 (0.87–2.04) (p = 0.1905)
<i>I usually feel rather lethargic</i>						
Not at all – little	183	40	21.9	Referent		
Somewhat – very much	97	27	27.8	1.27 (0.84–1.94)	1.25 (0.82–1.90)	1.34 (0.86–2.08) (p = 0.1939)
<i>I often feel exhausted</i>						
Not at all – little	179	38	21.2	Referent		
Somewhat – very much	101	29	28.7	1.35 (0.89–2.05)	1.34 (0.88–2.03)	1.52 (0.99–2.33) (p = 0.0541)
<i>I feel weary very much of the time</i>						
Not at all – little	193	46	23.8	Referent		
Somewhat – very much	87	21	24.1	1.01 (0.65–1.59)	0.99 (0.63–1.55)	1.13(0.72–1.79) (p = 0.5935)

^a Adjusted for age, gender, race/ethnicity, education, workload, physical activity, and alcohol consumption.

^b p-Value for the multivariable adjusted prevalence ratio.

helps to increase reaction time and alertness and improve performance. According to this idea, there should be a scheduled time for an officer to take a nap or an unrestricted opportunity to take a nap during the shift, as well as an appropriate place to take an uninterrupted nap. Police departments may wish to consider this napping strategy regardless of shift duration and include shifts less than 12 hours and those who might work double shifts.¹⁶ However, there is some controversy at organizational levels about napping since it takes away from patrol duties.

Mortality rates among police show significantly higher rates of cancers of the colon, kidney, digestive system, esophagus, male breast, and testis, as well as Hodgkin's disease.

Role of the Police Organization

Primary responsibility rests with the individual officer for getting proper sleep; however, police organizations can help their officers reach that goal. The organization can create a culture in which officers receive adequate information about the importance of good sleep habits, the hazards associated with fatigue and shift work, and strategies for managing them. According to Drs. Bryan Vila and Charles Samuels, organizations can do the following:

- During training at the basic academy level and in-service training afterward, educate personnel on the importance of proper sleep and the hazards associated with shift work.
- Fit the particular needs of each community and its police officers. There is no one ideal system for shift work.
- Shift scheduling strategies and staff deployment in police agencies require operational expertise and human factors expertise. Draw on these resources or perform research to guide shift development.
- Excessive overtime, frequent shift changes, and secondary employment are especially problematic issues in conjunction with sleep disorders and these elements should be carefully managed.¹⁷

Role of the Individual Officer

On an individual level, officers can improve the quality of their sleep by

- getting at least 7 hours of sleep daily;
- going to sleep at the same time every day, as much as possible;
- avoiding alcohol just before bedtime;
- using room-darkening curtains or shades (light keeps people awake);
- making their bedroom a place for sleep, not doing work or watching TV;
- maintaining a comfortable bedroom temperature—not too hot or cold;
- avoiding large meals or excessive fluids before bedtime;
- avoiding exercise within three hours of bedtime; and
- avoiding caffeine, nicotine, and other stimulants before bedtime.¹⁸

Conclusion

Shift work poses an increased negative risk among police officers in terms of safety and health. With proper sleep—and attention to this problem by both the organization and individual—such risks can be substantially reduced. It is likely that there will always be a

need for shift work in policing, so how law enforcement organizations adjust to this inherent work situation is essential to providing a high level of service to the community and to preserving the health and safety of officers. ♦

Notes:

¹Sleep Foundation, "Sleep in America Poll," 2010, <https://sleepfoundation.org/sites/default/files/nsaw/NSF%20Sleep%20in%20%20America%20Poll%20-%20Summary%20of%20Findings%20.pdf>.

²Bryan Vila and Dennis Kenney, "Tired Cops: The Prevalence and Potential Consequences of Police Fatigue," *NIJ Journal* 248 (2002): 16–21.

³Claudia Ma et al., "Associations of Objectively Measured and Self-Reported Sleep Duration with Carotid Artery Intima Thickness Among Police Officers," *American Journal of Industrial Medicine* 56 (2013): 1341–1351.

⁴Shantha M. W. Rajaratnam et al., "Sleep Disorders, Health, and Safety in Police Officers," *Journal of the American Medical Association* 306, no. 23 (2011): 2567–2578.

⁵William D. S. Killgore, "Effects of Sleep Deprivation on Cognition," *Progress in Brain Research* 185 (December 2010): 105–129.

⁶Timothy Roehrs et al., "Ethanol and Sleep Loss: A Dose Comparison of Impairing Effects," *Sleep* 26 (2003): 981–985.

⁷National Law Enforcement Officers Memorial Fund, *Preliminary 2017 Law Enforcement Officer Fatalities Report*, http://www.nleomf.org/assets/pdfs/reports/fatality-reports/2017/2017-End-of-Year-Officer-Fatalities-Report_FINAL.pdf.

⁸Destafekedulegn et al., "Fatigue and On-Duty Injuries Among Police Officers: The BCOPS Study," *Journal of Safety Research* 60 (February 2017): 43–51.

⁹Simon Folkard and Philip T. Tucker, "Shift Work, Safety and Productivity," *Occupational Medicine* 53, no. 2 (March 2003): 95–101.

¹⁰Gregory Belenky et al., "Patterns of Performance Degradation and Restoration During Sleep Restriction and Subsequent Recovery: A Sleep Dose-Response Study," *Journal of Sleep Research* 12, no. 1 (March 2003): 1–12.

¹¹John M. Violanti et al., "Life Expectancy in Police Officers: A Comparison with the U.S. General Population," *International Journal of Emergency Mental Health and Human Resilience* 15, no. 4 (2013): 217–228.

¹²Michael Wirth et al., "The Epidemiology of Cancer Among Police Officers," *American Journal of Industrial Medicine* 56, no. 11 (November 2013): 439–453.

¹³Ma et al., "Associations of Objectively Measured and Self-Reported Sleep Duration with Carotid Artery Intima Thickness Among Police Officers."

¹⁴Luenda E. Charles et al., "Shiftwork and Decline in Endothelial Function among Police Officers," *American Journal of Industrial Medicine* 59, no. 11 (November 2016): 1001–1008.

¹⁵John M. Violanti et al., "Atypical Work Hours and Metabolic Syndrome Among Police Officers," *Archives of Environmental and Occupational Health* 64, no. 3 (Fall 2009): 194–201.

¹⁶P. Daniel Patterson et al., "Evidence-Based Guidelines for Fatigue Risk Management in Emergency Medical Services," *Prehospital Emergency Care* 22, sup. 1 (2018): 89–101.

¹⁷Bryan Vila, Charles Samuels, and Nancy Jo Wesensten, "Sleep Loss in First Responders and the Military" in *Principles and Practice of Sleep Medicine*, eds. Meir H. Kryger and Thomas Roth, 6th ed. (Philadelphia, PA: Elsevier Saunders, 2016): 726–735.


¹⁸Michael J. Thorpy, "Managing the Patient with Shift-Work Disorder," *Journal of Family Practice* 59 (January 2010): S24–S31.

Supporting Officer Safety Through Family Wellness: The Effects of Sleep Deprivation

While sleep deprivation is not exclusive to law enforcement, it is often amplified due to the unique stressors of the job and shift work. Sleep is essential to maintain and repair bodily functions and systems. Sleep, or lack of, has effects on all functions of the mind and body, which not only affects an officer's job, but also family wellness.



Effects of Sleep Deprivation on the Body



Decreased Cognitive Processes, Problem-Solving, Concentration, and Reasoning

Lack of sleep inhibits decision making, interferes with forming sound judgements, and induces poor assessments due to increased irritability.ⁱ



Lack of Appetite Control

Sleep deprivation has shown to increase feelings of hunger and affect a person's ability to judge portion size.ⁱⁱ



Impaired Alertness

Sleep deprivation impairs an individual's ability to remain focused and alert for long periods of time.ⁱⁱⁱ



Increased Risk for Heart Disease

Sleep deprived persons are at a higher risk for heart disease and high blood pressure when only sleeping five to six hours a night.^{iv}



Spatial Disorientation

Sleep deprived persons are more likely to become disoriented when navigating, and often report slower reaction times.^v



More than half of all police officers don't get enough sleep.^{vi}

more than **40%** of police officers reported having **fallen asleep while driving**.

25% reported that happened **more than once a month**.^{vi}

Sleep deprivation can be dangerous



What about working the night shift?

Our bodies naturally relax and cool down when it gets dark outside and become alert when the sun is up. **Working second or third shift can disrupt the circadian clock and make sleep more difficult.** Working against the natural rhythms of the body can cause sleep disorders and fatigue.

Law enforcement is a 24-hour job and for many, working the night shift is unavoidable. Learning how to adapt to the schedule and demands of the job can help combat some of the potentially dangerous symptoms of working the night shift.

If you work a permanent night shift:

- It is best to slowly shift your circadian clock enough to still be able to function on days off.
- The best way to do this is on days off, go to sleep as late as possible and sleep as late as possible.
- On a workday, minimize the sunlight exposure on the drive home by wearing sunglasses and utilize blackout curtains when trying to sleep.^{ix}

If you work rotating night shifts:

- The circadian clock can't shift fast enough to keep up with a rotation.
- The American Psychological Association recommends avoiding symptomatic relief; caffeine to stay awake at night and sedatives to sleep during the day. These methods can be dangerous and only temporarily disrupt your circadian clock.^x



next page >

The Effects of Sleep Deprivation, continued...

What can officers do?

Practice a healthy lifestyle.

- Maintain balanced eating habits
- Refrain from tobacco use
- Limit alcohol consumption
- Exercise regularly
- Have an annual physical



Exercise proper sleep hygiene.

- Get seven to nine hours of sleep every night^{xi}
- Limit caffeine intake close to sleep time
- Minimize screen time before bed



Talk to your doctor about sleep disorders.

- Law enforcement officers are **twice as likely** to have a sleep disorder as non-law enforcement^{xii}



How can family help?

Assist in making the sleeping space **more comfortable and appealing**.

- Black out curtains, minimal electronics, supportive pillows and mattress, and a comfortable temperature are all ways to help with sleep.
- Ideal temperature for a room to sleep in is **60°-67°F**.^{xiii}
- Suggested addition: minimize activity in the house when the officer is trying to sleep.



Encourage your partner to talk to a doctor about his/her sleeping habits, particularly if s/he snores frequently, has trouble falling asleep or staying asleep, or begins to show other health concerns.

Create a family bedtime ritual.

Calming and relaxing environments help decrease stress and anxiety making falling asleep and staying asleep easier.

- One hour before you go to sleep have a 'wind down' hour. Do calming activities such as reading or taking a relaxing bath or shower. This is the crucial time to avoid electronic screens that can increase restlessness.^{xiv}



Get into a routine as much as possible.

Eating on a regular schedule and going to sleep and waking up on a regular schedule, no matter what the schedule is, all decrease the effects of sleep deprivation.

Communicate with family and friends to help distribute family responsibilities, such as sports practices, carpools, and grocery shopping.

Exercise regularly. Vigorous exercise can make it easier to fall asleep. Make it fun for the whole family. Think of creative family exercise opportunities like hiking, ice skating, dancing, and/or swimming.



- For more information about the Institute for Community-Police Relations, please visit: <http://www.theiacp.org/ICPR>.
- This publication is one in a series. For more family support resources please visit: <http://www.theiacp.org/ICPRlawenforcementfamily>

ⁱRajaratnam, Shantha MW, Laura K. Barger, Steven W. Lockley, Steven A. Shea, Wei Wang, Christopher P. Landrigan, Conor S. O'Brien et al. "Sleep disorders, health, and safety in police officers." *Jama* 306, no. 23 (2011): 2567-2578.

ⁱⁱHogenkamp, Pleunie S., Emil Nilsson, Victor C. Nilsson, Colin D. Chapman, Heike Vogel, Lina S. Lundberg, Sanaz Zarei et al. "Acute sleep deprivation increases portion size and affects food choice in young men." *Psychoneuroendocrinology* 38, no. 9 (2013): 1668-1674.

ⁱⁱⁱRajaratnam, Shantha MW, Laura K. Barger, Steven W. Lockley, Steven A. Shea, Wei Wang, Christopher P. Landrigan, Conor S. O'Brien et al. "Sleep disorders, health, and safety in police officers." *Jama* 306, no. 23 (2011): 2567-2578.

^{iv} <https://www.nhlbi.nih.gov/health/health-topics/topics/sdd/signs>. Updated June 2017.

^vValera, Silvana, Veronica Guadagni, Edward Slone, Ford Burles, Michele Ferrara, Tavis Campbell, and Giuseppe Iaria. "Poor sleep quality affects spatial orientation in virtual environments." *Sleep Science* 9, no. 3 (2016): 225-231.

^{vi}Rajaratnam, Shantha MW, Laura K. Barger, Steven W. Lockley, Steven A. Shea, Wei Wang, Christopher P. Landrigan, Conor S. O'Brien et al. "Sleep disorders, health, and safety in police officers." *Jama* 306, no. 23 (2011): 2567-2578.

^{vii}Rajaratnam, Shantha MW, Laura K. Barger, Steven W. Lockley, Steven A. Shea, Wei Wang, Christopher P. Landrigan, Conor S. O'Brien et al. "Sleep disorders, health, and safety in police officers." *Jama* 306, no. 23 (2011): 2567-2578.

^{viii}Dawson, Drew, and Kathryn Reid. "Fatigue, alcohol and performance impairment." *Nature* 388, no. 6639 (1997): 235.

^{ix} <http://www.apa.org/monitor/2011/01/night-work.aspx>. American Psychological Association, 2011.

^x <http://www.apa.org/monitor/2011/01/night-work.aspx>. American Psychological Association, 2011.

^{xi} National Sleep Foundation Memorandum released February 2015

^{xii} Pearsall, Beth. "Sleep disorders, work shifts and officer wellness." *National Institute of Justice Journal* 270 (2012): 36-39.

^{xiii} <https://sleep.org/articles/temperature-for-sleep/>

^{xiv} <https://sleepfoundation.org/sleep-tools-tips/healthy-sleep-tips/page/0/1>

This publication is one in a series. For more family support resources please visit: www.theiacp.org/ICPRlawenforcementfamily

The IACP's Institute for Community Police Relations (ICPR) is designed to provide guidance and assistance to law enforcement agencies looking to enhance community trust, by focusing on culture, policies, and practices.

For more information, contact us at...

@ ICPR@theIACP.org

theIACP.org/ICPR

800.THE.IACP

